

Babies and Toddlers

Meningitis Septicaemia Meningitis & Septicaemia



Meningitis and septicaemia often happen together. Be aware of all the signs and symptoms.

Symptoms can appear in any order. Some may not appear at all.

Early symptoms can include fever, headache, vomiting, muscle pain and fever with cold hands and feet.

Someone with meningitis or septicaemia can get a lot worse very quickly. Keep checking them.

**Trust your instincts –
get medical help immediately**

Children and Adults

Meningitis Septicaemia Meningitis & Septicaemia



Meningococcal Septicaemia

Meningococcal bacteria can cause meningitis and septicaemia (blood poisoning). One sign of septicaemia is a rash that does not fade under pressure – see Glass Test. Spots/rash may fade at first, keep checking.



- If someone is ill and getting worse do not wait for a rash, it can appear late or not at all.
- A fever with spots/rash that do not fade under pressure is a medical emergency.
- The spots or rash are more difficult to see on darker skin, check paler areas.

If you suspect meningitis or septicaemia

- Get medical help immediately. You can:
 - Call NHS Direct/NHS 24 or your GP
 - Go to your nearest accident and emergency department
 - Dial 999 for an ambulance
- Describe the symptoms and say you think it could be meningitis or septicaemia.
- Early diagnosis can be difficult. If you have had advice and are still worried, get medical help again.

**Trust your instincts –
get medical help immediately**



Common signs & symptoms of meningitis & septicaemia

The Meningitis Trust relies on voluntary donations to provide lifelong support for people affected by meningitis

helpline staffed by nurses
0800 028 18 28
www.meningitis-trust.org

Calls from BT landlines are free.
Other service providers and mobile rates may vary.

© Meningitis Trust June 2010
Registered charity number 803016 (England and Wales), SC037790 (Scotland)